

### Composition

Linseed meal, dextrose, dried carob pod meal, whey powder, wheat starch, yeasts (inactive saccharomyces cerevisiae), sodium chloride, sodium bicarbonate, psyllium seed kernel, banana flour, wheat flour, yeasts and parts thereof, cereals, colostrum, potassium chloride, calcium carbonate

### Analytical constituents and amounts

|                     |        |
|---------------------|--------|
| Crude protein ..... | 12.07% |
| Crude fat .....     | 2.80%  |
| Crude fibre .....   | 3.62%  |
| Crude ash .....     | 12.00% |
| Moisture .....      | 1.63%  |
| Sodium .....        | 2.75%  |
| Potassium .....     | 0.60%  |
| Chlorine .....      | 2.80%  |

### Additives per kg

#### Vitamins

|  |            |
|--|------------|
| 3a671 Vitamin D3 (cholecalciferol)                 | 12,000 IU  |
| 3a672a Vitamin A (retinyl acetate)                 | 100,000 IU |
| 3a700 Vitamin E (all rac-alpha-tocopheryl acetate) | 240 mg     |
| 3a821 Vitamin B1 (thiamine mononitrate)            | 16.00 mg   |
| 3a825i Vitamin B2 (riboflavine)                    | 12.00 mg   |
| 3a831 Vitamin B6 (pyridoxine hydrochloride)        | 8.00 mg    |
| Vitamin B12 (cyanocobalamin)                       | 160.20 mcg |

#### Trace elements

|  |         |
|--|---------|
| 3b202 Iodine (calcium iodate, anhydrous)             | 4.80 mg |
| 3b405 Copper (copper(II)sulphate, pentahydrate)      | 36 mg   |
| 3b503 Manganese (manganous(II)sulphate, monohydrate) | 240 mg  |
| 3b605 Zinc (zinc sulphate, monohydrate)              | 300 mg  |
| 3b801 Sodium selenite                                | 1.20 mg |
| 3b8.11 Selenised yeast inactivated NCYC R397         | 2.00 mg |



8 713296 021117



*it's about details*

# DiarFix

Dietetic supplementary feed  
for calves, for stabilisation  
of water and electrolyte  
balance to support the  
physiological digestion.

5 kg



### Instructions for use and dosage

Use Topro DiarFix in case of risk of, during periods of, or recovery from digestive upset (diarrhoea). It is recommended that a veterinarian's opinion be sought before use.

High dosage: 60 grams per litre of water for 1-3 days. Administer 2 liters to the calf twice a day inbetween the milk feeds.

Low dosage: 20 grams per litre of milk for 1-7 days.

### Storage instructions

Keep product cool, dry and out of reach of children. Protect against frost. Keep packaging closed after use.

### Manufacturing date

### Best before

### Batch No

### Registration No. manufacturer

α DE Ni 400036

GMP+ FSA assured  
GMO controlled  
PDV103322

Topro: registered trademark of



**bio enterprise**  
agro specialist

### Bio Enterprise B.V.

Sluiskolk 3, 7681 KC Vroomshoop NL  
info@topro.nl www.topro.nl